

# Between weak and Bruhat: the middle order on permutations


Mathilde Bouvel

*Loria, CNRS and Univ. Lorraine (Nancy, France).*


talk based on joint work with  
Luca Ferrari and Bridget E. Tenner

*Journées annuelles du GT Combinatoire Algébrique, Lyon, October 2024.*

# Familiar pictures




Bruhat order




weak order


# Familiar pictures and a third one



Bruhat order




middle order




weak order


# Familiar pictures and a third one



Bruhat order



middle order



weak order

## Goals of the talk:

- Define the middle order on  $S_n$
- Give meaning to the property that “it sits between weak and Bruhat”
- Describe some of its properties
- Popularize this new order, hoping you raise new questions about it


**Our 3 orders through “mesh patterns”**

# Permutation diagrams

## Notation:

- $S_n$  = the set of permutations of size  $n$
- $\sigma \in S_n$  is seen as the **word**  $\sigma(1)\sigma(2)\cdots\sigma(n)$
- $\sigma$  is also seen as its **diagram** i.e. the  $n \times n$  grid with points at coordinates  $(i, \sigma(i))$

**Example:** The word 1 8 3 6 4 2 5 7, or equivalently the diagram




# Permutation patterns

A permutation  $\pi \in S_k$  is a **pattern** of a permutation  $\sigma \in S_n$  (written  $\pi \preceq \sigma$ ) when there exist indices  $1 \leq i_1 < i_2 < \dots < i_k \leq n$  such that  $\sigma(i_a) < \sigma(i_b)$  if and only if  $\pi(a) < \pi(b)$


The subsequence  $\sigma(i_1)\sigma(i_2)\dots\sigma(i_k)$  is an **occurrence** of  $\pi$  in  $\sigma$

**Example:** 2134 is a pattern of 312854796, one occurrence being 3157

This is also seen on diagrams:



$\preceq$




# Permutation patterns

A permutation  $\pi \in S_k$  is a **pattern** of a permutation  $\sigma \in S_n$  (written  $\pi \preceq \sigma$ ) when there exist indices  $1 \leq i_1 < i_2 < \dots < i_k \leq n$  such that  $\sigma(i_a) < \sigma(i_b)$  if and only if  $\pi(a) < \pi(b)$


The subsequence  $\sigma(i_1)\sigma(i_2)\dots\sigma(i_k)$  is an **occurrence** of  $\pi$  in  $\sigma$

**Example:** 2134 is a pattern of 312854796, one occurrence being 3157

This is also seen on diagrams:



$\preceq$





# Permutation patterns

A permutation  $\pi \in S_k$  is a **pattern** of a permutation  $\sigma \in S_n$  (written  $\pi \preceq \sigma$ ) when there exist indices  $1 \leq i_1 < i_2 < \dots < i_k \leq n$  such that  $\sigma(i_a) < \sigma(i_b)$  if and only if  $\pi(a) < \pi(b)$

The subsequence  $\sigma(i_1)\sigma(i_2)\dots\sigma(i_k)$  is an **occurrence** of  $\pi$  in  $\sigma$

**Example:** 2134 is a pattern of 312854796, one occurrence being 3157

This is also seen on diagrams:

---

An **inversion** is a subsequence  $\dots j \dots i \dots$  in a permutation, with  $j > i$ .


Equivalently, it is an occurrence of the pattern 21 = 





**Example:** The inversions of 312854796 are  
31, 32, 85, 84, 87, 86, 54, 76 and 96

# Mesh patterns


A **mesh pattern**  $(\pi, M)$  is the data of a pattern  $\pi$  (say, of size  $k$ ) drawn in the central  $k \times k$  square of the grid  $[0, k + 1]^2$ , together with a set  $M$  of shaded unit cells in this grid. ( $M$  is called the mesh.)

An occurrence of  $(\pi, M)$  in  $\sigma$  is an occurrence of  $\pi$  in  $\sigma$  such that the regions of  $[0, n + 1]^2$  corresponding to the **mesh**  $M$  contain **no points** of  $\sigma$ .

**Example:** Consider the mesh pattern  $\mu =$ . The permutation 1423 contains four occurrences of 12, but only three of  $\mu$ .



# Weak order, seen through mesh patterns




Covering relations are described by

$$\dots ij \dots \rightsquigarrow \dots ji \dots$$

i.e., transforming an ascent<sup>(a)</sup> into a descent<sup>(d)</sup> using the same two values.

- (a) occurrence of 12 at consecutive positions
- (d) occurrence of 21 at consecutive positions

# Weak order, seen through mesh patterns




Covering relations are described by

$$\dots ij \dots \rightsquigarrow \dots ji \dots$$


i.e., transforming an ascent<sup>(a)</sup> into a descent<sup>(d)</sup> using the same two values.

- (a) occurrence of 12 at consecutive positions
- (d) occurrence of 21 at consecutive positions

Equivalently, covering relations are described by



# Bruhat order, seen through mesh patterns




**Relations** are described by the swaps

$$\dots i \dots j \dots \rightsquigarrow \dots j \dots i \dots$$

i.e., transforming a non-inversion (=occurrence of 12) into an inversion using the same two values.

**Covering relations** are the relations that do not create additional inversions.

# Bruhat order, seen through mesh patterns




**Relations** are described by the swaps

$$\dots i \dots j \dots \rightsquigarrow \dots j \dots i \dots$$

i.e., transforming a non-inversion (=occurrence of 12) into an inversion using the same two values.


**Covering relations** are the relations that do not create additional inversions.

**Equivalently**, covering relations are described by




# Middle order, defined through mesh patterns

- For the weak order, the covering relations are described by




- For the Bruhat order, the covering relations are described by




# Middle order, defined through mesh patterns


- For the weak order, the covering relations are described by



- For the middle order, the covering relations are described by




- For the Bruhat order, the covering relations are described by






# The middle order

- On permutations of size 3:




- Covering relations described by



# Summary so far, and what's ahead

## What we have seen:

- The covering relations of the middle order are described by




- This interpolates between the weak order and the Bruhat order

## What comes next:

# Summary so far, and what's ahead

## What we have seen:

- The covering relations of the middle order are described by



- This interpolates between the weak order and the Bruhat order

## What comes next:

- Another combinatorial interpretation of the middle order
- Some of its properties as a poset (in particular: distributive lattice)
- Enumeration of its intervals, and of its boolean intervals
- Implication on its Möbius function
- Combinatorial description of its Euler characteristic
- Restriction to the subset of involutions

# The middle order and inversion sequences

# Inversion sequences, and bijection with permutations

- Reminder: **Inversions** are occurrences  $\cdots j \cdots i \cdots$  of the pattern 21.
- $j$  is called **inversion top**.
- Given  $\sigma \in S_n$ , let  $x_j =$  number of inversions of  $\sigma$  such that  $j$  is the inversion top. Observe that  $0 \leq x_j < j$ .
- Let  $\varphi(\sigma) = (x_1, x_2, \dots, x_n)$  be the **inversion sequence** of  $\sigma$ .
- Sometimes called **Lehmer code**. Several (symmetric) variants exist.
- **Example:** For  $\sigma = 415623$ , we have  $\varphi(\sigma) = (0, 0, 0, 3, 2, 2)$
- This is a **bijection** between  $S_n$  and the set  $I_n$  of **inversion sequences** of size  $n$ :

$$I_n = [0, 0] \times [0, 1] \times [0, 2] \times \cdots \times [0, n-1]$$


# Middle order through inversion sequences

For inversion sequences  $\mathbf{x} = (x_1, x_2, \dots, x_n)$  and  $\mathbf{y} = (y_1, y_2, \dots, y_n)$ , define the **partial order**

$$\mathbf{x} \leq \mathbf{y} \text{ when } x_i \leq y_i \text{ for all } i$$


In particular, covering relations correspond to **adding 1 to one component** (provided we stay among inversion sequences).

**Theorem:** The middle order is the image of the above by the bijection  $\varphi^{-1}$ .




# Proving this characterization of the middle order



Let  $\varphi(\sigma) = (x_1, \dots, x_n)$  and  $\varphi(\tau) = (x_1, \dots, x_{j-1}, x_j + 1, x_{j+1}, \dots, x_n)$ .


- In particular,  $x_j < j - 1$ .
- So  $j$  is not a LtoR-minimum.
- So, we can define  $i$  as the rightmost entry to the left of  $j$  in  $\sigma$  such that  $i < j$ , and  $(i, j)$  is an occurrence of .
- We check that  $\tau$  is the permutation obtained swapping  $i$  and  $j$ , so that  $\tau$  covers  $\sigma$  in the middle order.

# Proving this characterization of the middle order

Let  $\varphi(\sigma) = (x_1, \dots, x_n)$  and  $\varphi(\tau) = (x_1, \dots, x_{j-1}, x_j + 1, x_{j+1}, \dots, x_n)$ .

- In particular,  $x_j < j - 1$ .
- So  $j$  is not a LtoR-minimum.
- So, we can define  $i$  as the rightmost entry to the left of  $j$  in  $\sigma$  such that  $i < j$ , and  $(i, j)$  is an occurrence of .
- We check that  $\tau$  is the permutation obtained swapping  $i$  and  $j$ , so that  $\tau$  covers  $\sigma$  in the middle order.

Let  $\tau$  be obtained from  $\sigma$  by transforming one  into .

- Let  $j$  be the largest of the two elements involved in .
- $\varphi(\sigma)$  and  $\varphi(\tau)$  differ only at their  $j$ -th coordinate
- and the difference is  $+1$
- meaning that  $\varphi(\tau)$  covers  $\varphi(\sigma)$  in the defined order on inversion sequences



## **First properties of the middle order**

# A product of chains

We have seen that the middle order  $\mathcal{P}_n$  is **isomorphic** (with explicit bijection  $\varphi$ ) **to the product of chains**

$$[0, 0] \times [0, 1] \times [0, 2] \times \cdots \times [0, n - 1]$$

Consequences:

- $\mathcal{P}_n$  is a **lattice**: any  $\sigma$  and  $\tau$  have a least upper bound  $\sigma \vee \tau$  (called **join**) and a greatest lower bound, denoted  $\sigma \wedge \tau$  (called **meet**).  
The join (resp. meet) is obtained taking **component-wise maximum (resp. minimum)** on corresponding inversion sequences.
- In addition,  $\mathcal{P}_n$  is a **distributive** lattice.  
(meaning that  $\vee$  is distributive over  $\wedge$  and vice-versa).
- $\mathcal{P}_n$  is **graded**, *i.e.* has a **rank function**  $r$ , meaning that, for any  $\sigma$ , we can define  $r(\sigma)$  as the length of any maximal chain from  $12 \cdots n$  to  $\sigma$ .  
In  $\mathcal{P}_n$ , we have  $r(\sigma) = \text{number of inversions of } \sigma$ .

**Intervals in the middle order**

# Characterizing and counting all intervals in $\mathcal{P}_n$

## Intervals of the $j$ -element chain $[0, j - 1]$ :

- Such intervals are
  - of the form  $\{a\}$  for  $0 \leq a \leq j - 1$ ,
  - or of the form  $[a, b]$  for  $0 \leq a < b \leq j - 1$ .
- Therefore, there are  $j + \binom{j}{2} = \binom{j+1}{2}$  such intervals.

## Intervals of $\mathcal{P}_n$ (up to isomorphism $\varphi$ ):

- Such intervals correspond to intervals  $[(x_1, \dots, x_n), (y_1, \dots, y_n)]$  where each  $[x_j, y_j]$  is an interval of  $[0, j - 1]$ .
- Therefore, there are  $\prod_{j=1}^n \binom{j+1}{2} = \frac{n!(n+1)!}{2^n}$  intervals in  $\mathcal{P}_n$ .

# Counting intervals by rank

**Fact:** An interval of rank  $k$  in  $\mathcal{P}_n$  corresponds to an interval  $[(x_1, \dots, x_n), (y_1, \dots, y_n)]$  with  $\sum x_j + k = \sum y_j$ .

**Theorem:** Denote by  $f(n, k)$  the number of intervals in  $\mathcal{P}_n$  having rank  $k$ , with  $n \geq 1$  and  $k \geq 0$ .

It holds that  $f(1, 0) = 1$  and for  $n \geq 2$  and  $k \in [0, \binom{n}{2}]$ ,

$$f(n, k) = \sum_{h=0}^{n-1} (n-h) \cdot f(n-1, k-h),$$

(with the convention that  $f(n, j) = 0$  when  $j < 0$ ).

**Proof:** By induction, decomposing an interval of rank  $k$  of  $[0, 0] \times [0, 1] \times \dots \times [0, n-2] \times [0, n-1]$  into an interval of rank  $h$  of  $[0, n-1]$  and an interval of rank  $k-h$  of  $[0, 0] \times [0, 1] \times \dots \times [0, n-2]$

# Table of $f(n, k)$ , which is A139769 on the OEIS

$n \backslash k$	0	1	2	3	4	5	6	7	8	9	10
1	1										
2	2	1									
3	6	7	4	1							
4	24	46	49	36	18	6	1				
5	120	326	501	562	497	354	204	94	33	8	1

- $k = 0$ :  $f(n, 0) = n!$  since it counts elements in  $\mathcal{P}_n$
- $k = 1$ :  $f(n, 1)$  counts the covering relations in  $\mathcal{P}_n$ . From the previous theorem, we have  $f(n, 1) = n!(n - H_n)$ , where  $H_n = \sum_{i=1}^n \frac{1}{i}$ .

This is sequence A067318 in the OEIS.

Another interpretation of this sequence is as **the sum of reflection lengths of all elements of  $S_n$** . Is there a bijective explanation?

**Boolean intervals in the middle order**

# Characterizing and counting boolean intervals in $\mathcal{P}_n$

**Dfn:** An interval is **boolean** if it is isomorphic to a boolean algebra.

**Characterization and enumeration:**

- Boolean intervals of  $\mathcal{P}_n$  correspond to pairs  $(x_1, \dots, x_n), (y_1, \dots, y_n)$  of inversion sequences with  $y_j \in \{x_j, x_j + 1\}$  for all  $j$ .
- The number of boolean intervals in  $\mathcal{P}_n$  is  $(2n - 1)!!$ .
  - Indeed, each pair  $(x_j, y_j)$  has  $j$  possibilities if  $y_j = x_j$ , and  $j - 1$  possibilities if  $y_j = x_j + 1$ , hence  $2j - 1$  possibilities.

**Properties about rank of boolean intervals:**

- The **rank** is the number of  $j$ 's such that  $y_j = x_j + 1$ .
- The maximal rank is  $n - 1$  (as  $x_1 = y_1 = 0$ ).



# Counting boolean intervals in $\mathcal{P}_n$ by rank

For  $n > k \geq 0$ , let  $b(n, k)$  be the number of rank- $k$  boolean intervals in  $\mathcal{P}_n$ .

- For  $n \geq j \geq 0$ , let  $c(n, j)$  be the number of permutations of size  $n$  having  $j$  cycles.
- Equivalently (using a variant of Foata's bijection),  $c(n, j)$  is the number of permutations of size  $n$  having  $j$  RtoL-minima.

**Theorem:**  $b(n, k) = \sum_{i=0}^n \binom{i}{k} c(n, n - i)$ .

**Proof:**

- $x_i = 0$  if and only if  $i$  is a RtoL-minimum. So  $c(n, n - i)$  is the number of inversion sequences of size  $n$  with  $i$  non-zero entries.
- A boolean interval of rank  $k$  in  $\mathcal{P}_n$  is characterized by an inversion sequence  $(y_1, \dots, y_n)$  with  $k$  non-zero entries marked (those such that  $x_j = y_j - 1$ ; we take  $x_j = y_j$  for non-marked entries).

# Möbius function

**Dfn:** The **Möbius function**  $\mu$  on any poset  $\mathcal{P}$  is defined recursively by

$$\mu(s, u) = \begin{cases} 0 & \text{if } s \not\leq u, \\ 1 & \text{if } s = u, \text{ and} \\ -\sum_{s \leq t < u} \mu(s, t) & \text{for all } s < u. \end{cases}$$

**Prop:** In finite distributive lattices, for any  $v, w$ , it holds that

- $\mu(v, w)$  is equal to 0 if the interval  $[v, w]$  is **not boolean**
- and otherwise  $\mu(v, w) = (-1)^t$ , where  $t$  is the rank of  $[v, w]$ .

This holds in  $\mathcal{P}_n$ . In particular, for  $\sigma \in S_n$ ,

- $\mu(12 \cdots n, \sigma) = 0$  if and only if  $\varphi(\sigma)$  contains  $\geq 1$  entry  $\geq 2$ .  
This is equivalent to  $\sigma$  containing a 312- or 321-pattern.
- Otherwise,  $\mu(12 \cdots n, \sigma) = (-1)^t$ , where  $t$  is the number of 1's in  $\varphi(\sigma)$  (here, also the number of inversions of  $\sigma$ ).

# Euler characteristic

# Euler characteristic of a finite distributive lattice

Let  $\mathcal{P}$  be a **finite distributive lattice**. (Recall that  $\mathcal{P}_n$  has this property.)

- **Dfn:** An element of  $\mathcal{P}$  is **join-irreducible** if it covers exactly one element of  $\mathcal{P}$ .
- **Dfn:** A **valuation** on  $\mathcal{P}$  is a function  $\nu$  that satisfies  $\nu(\min(\mathcal{P})) = 0$  and for all  $x, y$ ,

$$\nu(x) + \nu(y) = \nu(x \wedge y) + \nu(x \vee y).$$

- **Prop.:** A valuation is determined by its values on the join-irreducibles.
- **Dfn:** The **Euler characteristic** is the unique valuation  $\chi$  such that  $\chi(a) = 1$  for every join-irreducible  $a$ .

# Join-irreducibles and Euler characteristic in $\mathcal{P}_n$

**Prop.:** The **join-irreducible elements** of  $\mathcal{P}_n$  are the permutations

$$1 \ 2 \ \cdots \ i \ j \ (i+1) \ (i+2) \ \cdots \ (j-1) \ (j+1) \ \cdots \ n,$$

for  $i \in [0, n-2]$  and  $j \in [i+2, n]$ .

**Proof:** Similar to previous ones, using inversion sequences.

**Theorem:** The **Euler characteristic**  $\chi$  on  $\mathcal{P}_n$  is

$$\chi(\sigma) = \text{number of RtoL-non-minima of } \sigma.$$

**Proof:**

- $\chi$  is indeed 0 on  $12 \cdots n$ , and 1 on join-irreducibles.
- We can easily check that  $\chi(x) + \chi(y) = \chi(x \wedge y) + \chi(x \vee y)$ .



**Cor.:** The number of  $\sigma$  with Euler characteristic  $k$  in  $\mathcal{P}_n$  is  $c(n, n-k)$ .

**Proof:** As  $c(n, n-k) = \text{number of } \sigma \text{ of size } n \text{ with } k \text{ RtoL-non-minima}$

## **Restriction to involutions**

# Finding something not-so-nice in something too-beautiful

$\mathcal{P}_n$  is extremely well behaved. What about its restriction to [involutions](#)?



The subsequent poset  $\mathcal{I}_n$  is not a [lattice](#), not [graded](#), and not an [interval-closed](#) subposet of  $\mathcal{P}_n$ .

But ... we can still compute the [Möbius function](#) in  $\mathcal{I}_n$ .

# Characterizing inversion sequences of involutions

Recall that the number  $i(n)$  of involutions of size  $n$  satisfies

$$i(n) = i(n-1) + (n-1) \cdot i(n-2) \text{ for } n \geq 2.$$

The decomposition used to prove this recurrence also proves that:

**Prop.:** Let  $\sigma$  be a permutation and  $(x_1, \dots, x_n) = \varphi(\sigma)$  be its inversion sequence.

Then  $\sigma$  is an involution if and only if

- (i)  $x_n = 0$  and  $(x_1, \dots, x_{n-1})$  is the inversion sequence of an involution of size  $n-1$ , or
- (ii)  $x_n = k > 0$ ,  $x_{n-k} = 0$  and  $(x_1, \dots, x_{n-k-1}, x_{n-k+1} - 1, \dots, x_{n-1} - 1)$  is the inversion sequence of an involution of size  $n-2$ .



# Möbius function on $\mathcal{I}_n$


- We say that  $(x_1, \dots, x_n)$  is **slow-climbing** when it does not contain **large ascents**, defined as factors  $(x_i, x_{i+1})$  with  $x_{i+1} > x_i + 1$ .
- **Lemma:** The inversion sequence of an involution is slow-climbing if and only if it is the concatenation of factors  $(0, 1, \dots, h)$  for some (possibly different)  $h \geq 0$ .

**Theorem:** For any involution  $\sigma \in \mathcal{I}_n$ , let  $\alpha$  be the number of non-zero entries in  $\varphi(\sigma)$ . The Möbius function in  $\mathcal{I}_n$  is given by


$$\mu(12 \dots n, \sigma) = \begin{cases} (-1)^\alpha & \text{if } \sigma \text{ is slow-climbing, and} \\ 0 & \text{otherwise.} \end{cases}$$

**Wrapping up**


# Recap



Bruhat order



middle order




weak order


My goals were:

- Define the middle order on  $S_n$
- Give meaning to the property that “it sits between weak and Bruhat”
- Describe some of its properties
- Popularize this new order, hoping you raise new questions about it


# Recap



Bruhat order



middle order



weak order

My goals were:

- Define the middle order on  $S_n$
- Give meaning to the property that “it sits between weak and Bruhat”
- Describe some of its properties
- Popularize this new order, hoping you raise new questions about it

Thank you!